▶ Performance on the acute myocardial infarction mortality measure: **Hospitals with the highest proportions of vulnerable populations based on sociodemographic characteristics.** 

The Centers for Medicare & Medicaid Services (CMS) periodically investigates select hospital practices that may impact a hospital's performance on the following mortality measure: hospital-level 30-day risk-standardized mortality rate (RSMR) following acute myocardial infarction (AMI) [1]. The AMI mortality measure includes Medicare fee-for-service (FFS) and Veterans Health Administration (VA) beneficiaries aged 65 or older [2]. The AMI mortality measure assesses the occurrence of death for any cause within 30 days after hospital admission for AMI [2]. The AMI mortality measure has been publicly reported on <a href="Hospital Compare">Hospital Compare</a> since 2007 and has been included in the Hospital Value-Based Purchasing (HVBP) Program since 2013 [3].

There has been much discussion about the potential impact of patient sociodemographic status (SDS) on hospital outcome measures, including measures of mortality [3, 4]. We examined AMI RSMRs among hospitals identified as caring for a large proportion of vulnerable patients, as characterized by seven different SDS definitions, for hospitals with at least 25 eligible admissions. The hospitals included in this analysis are the 10% of hospitals with highest proportions of vulnerable patients as characterized by the different SDS definitions (Table 1). We compared the distribution of AMI RSMRs among the hospitals identified by the various SDS definitions and also examined the number of hospitals identified as caring for high proportion of vulnerable populations by the various definitions. Variation in RSMRs reflects differences in performance among hospitals; lower RSMRs suggest better quality, and higher RSMRs suggest worse quality. To ensure accurate assessment of each hospital, the AMI mortality measure uses a statistical model to adjust for key differences in patient risk factors that are clinically relevant and that have a strong relationship with the mortality outcome [2]. Please note that VA hospitals are not included in this analysis.

Among the subgroup of hospitals identified as serving vulnerable populations as characterized by the seven SDS definitions, only 47% of hospitals were identified by more than one definition of SDS.

Out of the 914 hospitals that met the criteria for serving a large proportion of vulnerable patients as characterized by at least one definition of SDS, less than half (428 hospitals; 47% of hospitals) met criteria characterized by multiple definitions of SDS. There were no hospitals that met criteria characterized by all seven definitions of SDS. However, eight hospitals (< 1% of 2,536 total hospitals) met the criteria for six of the definitions, 31 hospitals (1% of total hospitals) were identified by five of the definitions, 70 hospitals (3% of total hospitals) were identified by four of the definitions, and 105 hospitals (4% of total hospitals) were identified by three of the definitions.

The median AMI RSMR for all hospitals was 14.1% and median AMI RSMRs for hospitals identified as serving high proportions of vulnerable populations were within 0.4 percentage points of the overall median. We observed overlapping ranges in performance for hospitals meeting criteria based on all seven SDS definitions (Figure 1). The interquartile range of AMI RSMRs overall was 13.3% to 15.0% and for hospitals identified as serving high proportions of vulnerable populations the interquartile range was 13.4% to 15.2% (Table 2).







**TABLE I** Definitions and data sources for the following sociodemographic characteristics: living below the U.S. poverty line; an educational attainment below high school; unemployed; living in crowded households; African-American; receiving Medicaid; and residing in a zip-code with a low AHRQ Index of SES score.


Sociodemographic characteristics	Definition of vulnerable patients based on sociodemographic characteristics	Identification of hospitals with the highest proportion of vulnerable patients	Source
Below U.S. poverty line	Patients from zip codes where more than 29.7% of the residents are below the United States (U.S.) poverty line	Hospitals with more than 56.7% of Medicare fee-for service (FFS) patients that meet the vulnerable patient definition	American Community Survey (ACS) 2008 – 2012 5 year estimate Medicare Part A Inpatient Claims 2012
Educational attainment below high school	Patients from zip codes where more than 18.6% of the residents aged ≥ 25 years have less than a 12th-grade education	Hospitals with more than 69.7% of Medicare FFS patients that meet the vulnerable patient definition	ACS 2008 – 2012 5 year estimate Medicare Part A Inpatient Claims 2012
Unemployed	Patients from zip codes where more than 11.5% of the residents aged 16 years or older in labor force who are unemployed and actively seeking work	Hospitals with more than 64.4% of Medicare FFS patients that meet the vulnerable patient definition	ACS 2008 – 2012 5 year estimate Medicare Part A Inpatient Claims 2012
Crowded households	Patients from zip codes where more than 3.2% of the residents live in households containing one or more person per room	Hospitals with more than 70.2% of Medicare FFS patients that meet the vulnerable patient definition	ACS 2008 – 2012 5 year estimate Medicare Part A Inpatient Claims 2012
African-American	African-American patients	Hospitals with more than 23.7% of Medicare FFS patients identified as African-American	Medicare Part A Inpatient Claims 2013
Medicaid	Patients that have Medicaid coverage	Hospitals with more than 30.5% of patients with Medicaid coverage	American Hospital Association (AHA) Survey 2013 [5]
AHRQ Index of SES scores	Patients from zip codes with an Agency for Healthcare Research & Quality (AHRQ) socioeconomic status (SES) index score below 31.8	Hospitals with more than 72.7% of Medicare FFS patients that meet the vulnerable patient definition	AHRQ SES index [6] ACS 2008 – 2012 5 year estimate Medicare Part A Inpatient Claims 2012







**FIGURE I** Distributions and medians of AMI RSMRs (%) for hospitals with the highest proportion of patients with one of the following sociodemographic characteristics: living below the U.S. poverty line; an educational attainment below high school; unemployed; living in crowded households; African-American; receiving Medicaid; and residing in a zip-code with a low AHRQ Index of SES score, July 2011-June 2014.









**TABLE 2** Distributions of AMI RSMRs (%) for hospitals with the highest proportion of patients with one of the following sociodemographic characteristics: living below the U.S. poverty line; an educational attainment below high school; unemployed; living in crowded households; African-American; receiving Medicaid; and residing in a zip-code with a low AHRQ Index of SES score, July 2011-June 2014.

## AMI RSMRs (%) for hospitals with the highest proportions of patients:

	All hospitals; n=2,536	Below U.S. poverty line; n=245	Educational attainment below high school; n=245	Unemployed; n=245	In crowded households; n=244	African- American; n=245	Medicaid; n=243	Low AHRQ Index of SES score; n=245
Maximum	20.6	18.6	17.8	18.2	17.5	17.9	18.1	17.8
75%	15.0	15.0	15.1	15.1	14.8	15.1	15.1	15.2
Median (50%)	14.1	14.0	14.4	14.3	14.1	14.2	14.3	14.5
25%	13.3	13.4	13.6	13.6	13.4	13.4	13.5	13.6
Minimum	9.9	10.4	10.4	10.4	10.4	10.4	11.0	11.5

The median AMI RSMR for all hospitals was 14.1% and median AMI RSMRs for hospitals identified as serving high proportions of vulnerable populations were within 0.4 percentage points of the overall median. Similarly, the interquartile range of AMI RSMRs overall was 13.3% to 15.0% and for hospitals identified as serving high proportions of vulnerable populations the interquartile range was 13.4% to 15.2%.

- 1. Medicare Hospital Quality Chartbook 2014: Performance Report on Outcome Measures. Prepared by Yale New Haven Health Services Corporation Center for Outcomes Research and Evaluation for the Centers for Medicare and Medicaid Services 2014; <a href="http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/Downloads/Medicare-Hospital-Quality-Chartbook-2014.pdf">http://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Assessment-Instruments/HospitalQuality-Initiatives-Patient-Instruments/HospitalQuality-Initiatives-Patient-Instruments/HospitalQuality-Initiatives-Patient-Instruments/HospitalQuality-Initiatives-Patient-Instruments/HospitalQuality-Initiatives-Patient-Instruments/HospitalQuality-Initiatives-Instruments/HospitalQuality-Initiatives-Initiatives-In
- 2. Dorsey K, Grady J, Desai N, et al. 2015 Condition-Specific Measures Updates and Specifications Report Hospital-Level 30-Day Risk-Standardized Mortality Measures: Acute Myocardial Infarction Version 9.0, Heart Failure Version 9.0, Pneumonia Version 9.0, Chronic Obstructive Pulmonary Disease Version 4.0, Stroke Version 4.0; https://www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic%2FPage%2FQnetTier4&cid=1163010421830. Accessed 26 June 2015.
- 3. "Hospital Inpatient Prospective Payment Systems for Acute Care Hospitals, Final Rule." Federal Register / 22 August 2014; http://federalregister.gov/a/2014-18545. Accessed 16 June 2015.
- 4. National Quality Forum. Risk Adjustment for Socioeconomic Status or Other Sociodemographic Factors, Technical Report, August 15, 2014; <a href="http://www.qualityforum.org/Publications/2014/08/Risk Adjustment for Socioeconomic Status or Other Sociodemographic Factors.aspx">http://www.qualityforum.org/Publications/2014/08/Risk Adjustment for Socioeconomic Status or Other Sociodemographic Factors.aspx</a>. Accessed 9 July 2015.
- 5. AHA Annual Survey Database Fiscal Year 2013; http://www.ahadataviewer.com/book-cd-products/aha-survey/. Accessed 26 June 2015.
- 6. U.S. Department of Health & Human Services, AHRQ Agency for Healthcare Research and Quality archive; Publication # 08-0029-EF, Chapter 3: Creation of New Race-Ethnicity Codes and SES Indicators for Medicare Beneficiaries Chapter 3: Creating and Validating and Index of Socioeconomic Status; <a href="http://archive.ahrq.gov/research/findings/final-reports/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicareindicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicators/medicat





